

MADPH-99-1146

hep-ph/9911442

Top-quark decay via R -parity violating interactions at the Tevatron

T. Han and M. B. Magro

Department of Physics, University of Wisconsin, Madison, WI 53706, USA

(January, 2000)

Abstract

We consider the top-quark decay $t \rightarrow \tilde{\tau} b$ and $t \rightarrow \tau b \tilde{\chi}_1^0$ via explicit R -parity violating interactions in SUSY theories. We discuss the observability of those channels at the Fermilab Tevatron collider. The existing Tevatron data indicate a 95% confidence level upper bound on the coupling to be 0.94 (0.63) for a long-lived (short-lived) $\tilde{\chi}_1^0$ with $m_{\tilde{\tau}} = 70$ GeV. At Tevatron Run II with an integrated luminosity of 2 (10) fb^{-1} , one can obtain a 2σ constraint as 0.38 (0.24) for a long-lived $\tilde{\chi}_1^0$ and 0.29 (0.19) for a short-lived $\tilde{\chi}_1^0$, beyond the current indirect limit.

11.30.Fs, 12.60.Jv, 13.90.+i, 14.65.Ha

Typeset using REVTeX

I. INTRODUCTION

Physics associated with the top quark is fascinating. Being the heaviest elementary particle observed so far with a mass $m_t \approx 175$ GeV, the top quark has the greatest potential to probe new physics beyond the Standard Model (SM). Examples considered include the radiative electroweak symmetry breaking due to the large top-quark Yukawa coupling in supersymmetric (SUSY) grand unified theories (GUT), and the dynamical electroweak symmetry breaking in the top-quark sector. Furthermore, the fact that the top quark is about 35 times more massive than its SU(2)-doublet partner, the b quark, makes it an attractive candidate for studying flavor physics.

The Fermilab Tevatron will start a new run with significant upgrade on luminosity, reaching approximately 2 fb^{-1} in a year. This would yield more than ten thousand $t\bar{t}$ pairs and thus provide a great opportunity to explore new physics in the top-quark sector. Observing the top-quark rare decays may reveal information on new physics. For instance, if there are charged scalars such as the charged Higgs bosons ($H^\pm$), the decay $t \rightarrow H^+ b$ may occur if kinematically accessible. This search has been carried out with the existing data from the Tevatron by the CDF and D0 collaborations [1,2]. The non-observation of the predicted signal has established limits on the charged Higgs boson mass and its coupling to the top quark. In SUSY GUT models, one may be motivated to search for decays to scalar quarks and gauginos [3]

$$t \rightarrow \tilde{t}_1 \tilde{\chi}_1^0 \quad \text{and} \quad \tilde{b} \tilde{\chi}_1^+ \quad (1)$$

where $\tilde{t}_1$ is the lighter top-squark and $\tilde{\chi}_1^0$ the lightest neutralino, assumed to be the lightest supersymmetric particle. Furthermore, if R -parity (R_P) is broken [4,5], then $\tilde{\chi}_1^0$ will not be stable and even possibly decay within the detector, leading to a different signature in the Tevatron collider experiments [6,7].

Process Eq. (1) may be disfavored by kinematics for heavy SUSY particles. Also, generic R_P -breaking ($\mathcal{R}_P$) interactions are subject to stringent experimental constraints due to the

absence of proton decay and large flavor-changing neutral currents in the lepton and quark sectors [5]. However, the bounds on operators involving the third generation fermions are particularly weak, which reflects our lack of knowledge for flavor physics in the third generation. If there exist sizeable $\mathcal{R}_p$ interactions involving the third generation, then the top quark may directly decay to other interesting channels. In fact, decays of the top quark and the top squark have been studied in scenarios where R_P is spontaneously broken [8]. In this Letter, we explore the explicit $\mathcal{R}_p$ couplings

$$\begin{aligned} & \lambda'_{333} (\overline{\nu_{\tau L}^c} b_L \tilde{b}_R^* + \bar{b}_R b_L \tilde{\nu}_{\tau L} + \bar{b}_R \nu_{\tau L} \tilde{b}_L \\ & - \overline{\tau_L^c} t_L \tilde{b}_R^* - \bar{b}_R t_L \tilde{\tau}_L - \bar{b}_R \tau_L \tilde{t}_L) + h.c. \end{aligned} \quad (2)$$

where the subscript for the coupling indicates the third generation for all particles involved, and τ_L^c is the charge conjugate field. The above interactions lead to the top-quark decay to

$$t \rightarrow b \tilde{\tau}^+ \rightarrow b \tau^+ \tilde{\chi}_1^0, \quad (3)$$

$$t \rightarrow \tilde{b} \tau^+ \rightarrow b \tilde{\chi}_1^0 \tau^+, \quad (4)$$

$$\text{and } t \rightarrow \tilde{t}_1 \tilde{\chi}_1^0 \rightarrow b \tau^+ \tilde{\chi}_1^0. \quad (5)$$

Motivated by SUSY GUT models, we assume that $\tilde{\tau}$ is lighter than the top quark and it decays dominantly to $\tau \tilde{\chi}_1^0$, as indicated in Eq. (3). $\tilde{b}$ is heavier and the $t \rightarrow \tilde{b} \tau^+$ channel in Eq. (4) is a three-body decay via a virtual $\tilde{b}^*$. We also take $m_{\tilde{t}_1} + m_{\tilde{\chi}_1^0} > m_t$ so that the channels of Eq. (1) are kinematically suppressed as we only have the three-body decay shown in Eq. (5) via a virtual $\tilde{t}_1^*$.

There exist weak indirect bounds on the coupling λ'_{333} . From the $Z \rightarrow \tau^+ \tau^-$ partial width via the $\mathcal{R}_p$ radiative corrections [9,10], a 2σ limit $|\lambda'_{333}| < 0.58$ was obtained for $m_{\tilde{b}} = 100$ GeV. Considering the decay rate of $B \rightarrow \tau \nu_\tau X$ yielded a bound of similar size [11]. The bounds become weaker for a heavier $\tilde{b}$. To concentrate on this possibly large coupling, we assume that the other $\mathcal{R}_p$ couplings involving light fermions are much smaller. The collider phenomenology then depends on whether $\tilde{\chi}_1^0$ in Eqs. (3) to (5) would decay within the detector or not, as we will discuss in details.




FIG. 1. Feynman diagrams for top-quark two-body (a) and three-body (b,c) decays via R -parity violation operators in Eq. (2).

The rest of the paper is organized as follows: In Sec. II, we discuss the two-body and three-body top decay widths due to λ'_{333} and obtain the respective branching fractions. In Sec. III, we study the new signatures coming from these decays and the sensitivity to probe λ'_{333} at the Tevatron collider. We make some concluding remarks in Sec. IV.

II. TOP DECAYS

We evaluate the partial decay widths for processes (3), (4) and (5) depicted in Fig. 1. Denoting the two- and three-body decay widths by Γ_2 and Γ_3 , and neglecting the masses of b and τ , we obtain for the two-body decay

$$\Gamma_2(t \rightarrow b\tilde{\tau}) = \frac{|\lambda'_{333}|^2}{32\pi} m_t \left(1 - \frac{m_{\tilde{\tau}}^2}{m_t^2}\right)^2, \quad (6)$$

which agrees with Ref. [6]; and for the three-body decay

$$\Gamma_3(t \rightarrow \tau b\tilde{\chi}_1^0) = \frac{|\lambda'_{333}|^2}{3072\pi^3} \left((C_L^2 + C_R^2) \frac{m_t^5}{m_b^4} I_b(x) + (C_L'^2 + C_R'^2) \frac{m_t^5}{m_t^4} I_t(x) \right), \quad (7)$$

with $C_{L,R}$ and $C'_{L,R}$ being the left- and right-handed coupling constants for the vertices $b\tilde{b}\tilde{\chi}_1^0$ and $t\tilde{t}\tilde{\chi}_1^0$ respectively [12]. The phase space integrals $I_i(x)$ are given by

$$I_b(x) = 12 \int_{x_\chi}^1 dx \frac{(1-x)^2(x-x_\chi)^2}{x(1-x/x_b)^2}, \quad (8)$$

and

$$I_{\bar{t}}(x) = 12 \int_0^{(1-x_\chi^{1/2})^2} dx \frac{x}{(1-x/x_{\bar{t}})^2} \left[(1+x_\chi-x) + 4x_\chi^{1/2}A \right] \lambda^{1/2}(1, x_\chi, x), \quad (9)$$

where $x_i = m_i^2/m_t^2$, $A = C'_L C'_R / (C'^2_L + C'^2_R)$ and λ is the usual Källén kinematical function. The factor of 12 is to normalize the integrals $I_{\bar{b}}$ and $I_{\bar{t}}$ to unity in the limit $x_{\tilde{\chi}_1^0} \rightarrow 0$ and $x_{\bar{b}}, x_{\bar{t}} \gg 1$. The ratio of these two decay widths is $\Gamma_2/\Gamma_3 \approx 48\pi^2(C_L^2 + C_R^2)m_t^4/m_{\bar{t}}^4$, which shows that the two-body mode dominates by more than three orders of magnitude.

The SM partial width for the top-quark decay $t \rightarrow bW^+$ is

$$\Gamma_{SM}(t \rightarrow bW^+) = \frac{g^2 m_t^3}{64\pi M_W^2} |V_{tb}|^2 \left(1 - \frac{m_W^2}{m_t^2}\right) \left(1 + \frac{m_W^2}{m_t^2} - 2\frac{m_W^4}{m_t^4}\right). \quad (10)$$

The ratio for the $\cancel{R}_p$ decay width with respect to that of the SM is thus

$$\Gamma_2/\Gamma_{SM} \approx 2|\lambda'_{333}|^2 M_W^2/g^2 m_t^2 \approx |\lambda'_{333}|^2. \quad (11)$$

We see that a coupling $|\lambda'_{333}| \sim 0.5$ would allow a branching ratio of about 20%.

In Fig. 2 we show the magnitude for the branching fractions of the two-body decay for $m_{\tilde{\tau}} = 70$ and 100 GeV with $m_{\tilde{\chi}_1^0} = 60$ GeV and three-body decay for $m_{\tilde{\chi}_1^0} = 60$ and 80 GeV with $m_{\tilde{\tau}} = 70$ GeV, as a function of the coupling constant $|\lambda'_{333}|$. We have chosen $m_{\bar{b}} = m_{\bar{t}_1} = 200$ GeV and $(C_L^2 + C_R^2) \approx (C'^2_L + C'^2_R) \approx 0.1$, with $A \ll 0.1$, which enhances the three-body decay and is compatible with the masses we have assumed for the lightest neutralino. As one can see, the two-body decay can be very important reaching 10 – 20% or higher for large $|\lambda'_{333}|$ and therefore could play an important role in the top decay phenomenology. On the other hand, the three-body decay is too small to be significant. For illustration, we have also included the SM channel ($t \rightarrow W^+ b$) in Fig. 2 at the top with $m_{\tilde{\tau}} = 70$ GeV.

III. SIGNATURES

We base our analyses on the SM top-quark production at the Tevatron

$$p\bar{p} \rightarrow t\bar{t}X, \quad (12)$$




FIG. 2. R -parity violating branching fractions for top decays as a function of the coupling constant $|\lambda'_{333}|$. The two upper parallel curves represent the two-body decay $t \rightarrow b\tilde{\tau}^+$, for $m_{\tilde{\tau}} = 70$ GeV (dashed) and $m_{\tilde{\tau}} = 100$ GeV (solid) with $m_{\tilde{\chi}_1^0} = 60$ GeV; while the two lower parallel curves stand for the three-body decay, $t \rightarrow b\tau^+\tilde{\chi}_1^0$, for $m_{\tilde{\chi}_1^0} = 60$ GeV (dashed) and $m_{\tilde{\chi}_1^0} = 80$ GeV (solid) with $m_{\tilde{\tau}} = 70$ GeV. We assumed $m_{\tilde{b}} = m_{\tilde{t}_1} = 200$ GeV for the three-body decay. The dot-dashed curve at the top is for the SM channel ($t \rightarrow W^+b$) with $m_{\tilde{\tau}} = 70$ GeV.

from those recently reported in the CDF/D0 experiments [13]. The $t\bar{t}$ signal can be identified above the SM backgrounds from their distinctive decay products. In our analyses, the event acceptance can be expressed by

$$\mathcal{A} = \sum_{i,j}^{SM, \cancel{R}_p} \epsilon_{ij} B_i B_j, \quad (13)$$

where B_{SM} and $B_{\cancel{R}_p}$ are the branching fractions of t and $\bar{t}$ decays to $t \rightarrow bW$ and $t \rightarrow b\tau\tilde{\chi}_1^0$, ϵ_{ij} the corresponding detection efficiencies.

With the $\cancel{R}_p$ interactions under consideration, the LSP $\tilde{\chi}_1^0$ is not stable and will decay via $\tilde{\chi}_1^0 \rightarrow \nu_\tau b\bar{b}$. The collider phenomenology will then depend upon if it decays within the detector or not. The decay length is calculated by $\ell = \gamma\beta c\tau$, where the $\tilde{\chi}_1^0$ life-time τ can be written as [5]




FIG. 3. Estimated decay length for $\tilde{\chi}_1^0 \rightarrow \nu_\tau b \bar{b}$ with various effective coupling κ , assuming $\gamma\beta = 3$. The dashed line indicates the five-meter mark.

$$\tau^{-1} = \kappa^2 \frac{3 G_F^2 m_{\tilde{\chi}_1^0}^5}{192 \pi^3}, \quad (14)$$

where the effective coupling κ depends on the composition of the LSP and the masses of the virtual sfermions [5,14], and is typically $\kappa \sim 0.1 (100 \text{ GeV}/m_{\tilde{f}})^2 \lambda'_{333}$. At the Tevatron energies, the $\gamma\beta$ factor for $\tilde{\chi}_1^0$ is typically about $1 \sim 5$. With this estimation, we can calculate the $\tilde{\chi}_1^0$ decay length as shown in Fig. 3. We see that for a rather light LSP $m_{\tilde{\chi}_1^0} \sim \mathcal{O}(10 \text{ GeV})$ or a weak effective coupling $\kappa < 10^{-5}$, $\tilde{\chi}_1^0$ may escape from detection before it decays, resulting in missing energy as the standard SUSY signal. In fact, there is no mass constraint of this order for the LSP for general SUSY models. Also, such a small coupling is possible even with $\lambda'_{333} \sim \mathcal{O}(0.1 - 1)$ if the third generation sfermions are very massive about 10 TeV as advocated in certain “inverse hierarchy” models [15]. On the other hand, for a large region of the parameter space in $(\kappa, m_{\tilde{\chi}_1^0})$, $\tilde{\chi}_1^0$ will decay within the detector of a few meters. We will therefore study both cases below.

A. Long-lived $\tilde{\chi}_1^0$ as missing transverse energy




FIG. 4. Predicted number of events versus the coupling $|\lambda'_{333}|$ for Run I ($\sqrt{s} = 1.8$ TeV) with the D0 event reconstruction method and a data sample of 109.2 pb^{-1} for two representative values of $m_{\tilde{\tau}}$. The solid curves stand for a long-lived $\tilde{\chi}_1^0$ and the dashed ones for $\tilde{\chi}_1^0$ to decay within detector.

We consider the simple situation first: The LSP $\tilde{\chi}_1^0$ is stable in the experimental scale, resulting in missing transverse energy. It is important to note that in this case the top-quark decay via the $\mathcal{R}_p$ coupling λ'_{333} yields the same signal as for the charged Higgs boson search $H^+ \rightarrow \tau^+ \nu b$. The efficiencies for these final states are given by [2]

$\epsilon_{ij} \text{ (%)}$	$t \rightarrow bW$	$t \rightarrow b\tau\nu$	
$t \rightarrow bW$	3.42	1.36	(15)
$t \rightarrow b\tau\nu$	1.36	0.41	

We adopt the D0 analyses and assume the same efficiencies for the event reconstruction. The predicted event rate can thus be obtained by

$$N = \mathcal{A} \sigma(t\bar{t}) \mathcal{L} + N_B, \quad (16)$$

where $\sigma(t\bar{t}) \approx 5.5$ pb [2], $\mathcal{L}$ is the integrated luminosity. N_B presents the expected non-top-quark SM backgrounds, which includes W +jets events and QCD multi-jet events with a misidentified lepton and large $\cancel{E}_T$ [2]. Based on this event reconstruction scheme, we calculate the predicted $t\bar{t}$ signal events as shown by the solid curves in Fig. 4. We have included all the four channels listed in Eq. (15). The SM expectation for $t\bar{t}$ events is implied at $|\lambda'_{333}| = 0$. The larger the $\mathcal{R}_p$ coupling is, the smaller the SM branching fraction (B_{SM}) would be, as seen from the top curve in Fig. 2. Since the detection efficiencies are optimized for the SM decay channels, the total predicted number of $t\bar{t}$ events decreases for higher values of $|\lambda'_{333}|$ after convoluting with the efficiencies of Eq. (15), resulting in a “disappearance” experiment.

The non-observation of the signal beyond the SM at the Tevatron experiments can be translated to constraints on $m_{\tilde{\tau}}$ and $|\lambda'_{333}|$. In Fig. 5(a), we show the exclusion limit in the $m_{\tilde{\tau}} - |\lambda'_{333}|$ plane at the 95% confidence level. Here we have adopted the Poisson statistics treatment because the number of signal events with the existing data is not expected to be very large. Since our procedure closely follows the D0 analysis [2], we assume the D0 data sample of 109.2 pb^{-1} and obtain the solid curve. The dashed curve is for 200 pb^{-1} (CDF and D0). The regions below the curves are excluded at a 95% C.L. level. For the current mass limit $m_{\tilde{\tau}} \approx 70 \text{ GeV}$ [16], we see that the existing Tevatron data can put a bound on the coupling to be $|\lambda'_{333}| < 0.94$. The bound becomes weaker for a heavier $m_{\tilde{\tau}}$. At Run II with $\sqrt{s} = 2 \text{ TeV}$ and $\sigma(t\bar{t}) \approx 6.8 \text{ pb}$, the sensitivity is expected to be greatly enhanced. This is shown in Fig. 6(a). For $m_{\tilde{\tau}} = 70 \text{ GeV}$, the 2σ bound is $|\lambda'_{333}| < 0.38$ (0.24) with 2 (10) fb^{-1} . This bound is essentially independent of the other SUSY parameters, and is more stringent than the indirect one obtained in Ref. [9,10]. For comparison, the 3σ contours are also given by the dashed curves in the figure.




FIG. 5. Contour plots for Run I in the $m_{\tilde{\tau}} - |\lambda'_{333}|$ plane for exclusion at the 95% confidence level for the cases where (a) the $\tilde{\chi}_1^0$ is long-lived and (b) the $\tilde{\chi}_1^0$ decays within the detector. The solid curves are for the data sample of 109.2 pb^{-1} (D0), and the dashed curves are for 200 pb^{-1} .




FIG. 6. Contour plots in the $m_{\tilde{\tau}} - |\lambda'_{333}|$ plane for exclusion at the 2σ (solid) and 3σ (dashed) limits at Run II with 2 and 10 fb^{-1} , for the cases where (a) the $\tilde{\chi}_1^0$ is long-lived and (b) the $\tilde{\chi}_1^0$ decays within the detector.

B. $\tilde{\chi}_1^0 \rightarrow \nu_\tau b \bar{b}$

As seen in Fig. 3, if $m_{\tilde{\chi}_1^0}$ and κ are large $\tilde{\chi}_1^0$ can decay within detector, yielding extra $b\bar{b}$ plus missing transverse energy in the final state. Two more b quarks would make the signal very different from the SM mode. This decay was considered at the Tevatron in the light of the total $t\bar{t}$ production cross section and a rather loose bound was inferred [14,11]. We here consider two searching scenarios.

First, we still adopt the disappearance method because the study of SM $t\bar{t}$ events is well established. Since the signal mode

$$t \rightarrow \tau^+ b \tilde{\chi}_1^0 \rightarrow \tau^+ b b \bar{b} \nu_\tau \quad (17)$$

is significantly different from the mode $t \rightarrow \tau b \nu_\tau$ being searched for, the signal events can be effectively removed from the sample, leading to a lower acceptance efficiency. We thus keep the SM $t \rightarrow bW$ mode with the same efficiency as in Eq. (15), while take the others to be zero. This is an optimistic choice for the signal searches in a disappearance experiment, but it will not significantly alter the results. The “more disappearance” results in a larger deviation from the SM expectation of $t\bar{t}$ production, as shown by the dashed curves in Fig. 4. This in turn leads to more stringent constraints with the existing Tevatron data as shown in Fig. 5(b) and with Run II data in Fig. 6(b). The current Tevatron data can put a bound on the coupling to be about $|\lambda'_{333}| < 0.63$ for the short-lived $\tilde{\chi}_1^0$ with $m_{\tilde{\tau}} \sim 70$ GeV. At Run II, the 2σ bound will be $|\lambda'_{333}| < 0.29$ (0.19) with 2 (10) fb^{-1} .

Alternatively, we could consider the direct search for the signal channel Eq. (17). To have a clean signal, one leptonic decay of $t \rightarrow b\ell\nu$ ($\ell = e, \mu$) is needed in combination with the signal mode. This $\ell\tau$ channel has been recently studied experimentally in Ref. [17]. Unfortunately, the low branching fraction for the leptonic modes and the low detection efficiency for the τ lepton render the signal events too low to be appreciable. Only with a luminosity significantly higher than the Run II projection could this search become possibly useful. We note that direct production processes $p\bar{p} \rightarrow t\bar{\tau}X$ have been suggested to search for this $\mathcal{R}_p$ interaction [18] and it is worthwhile exploring further.

IV. DISCUSSIONS AND CONCLUSION

Our analyses closely follow the study reported in Ref. [2] for $H^\pm \rightarrow \tau^\pm \nu_\tau$; while our signal for a long-lived $\tilde{\chi}_1^0$ is for $\tilde{\tau}^\pm \rightarrow \tau^\pm \tilde{\chi}_1^0$. Although one may expect to have a softer τ lepton in our signal than that in Ref. [2] due to a more massive $\tilde{\chi}_1^0$, we would like to argue that the event reconstruction efficiency may be quite comparable since the signal under consideration carries larger missing energy, which should help in event selection.

A more interesting question to ask would be that if there is a $\tau^\pm$ +missing energy signal established beyond the SM expectation, could we tell if it is from a $H^\pm$ decay or from $\tilde{\tau}^\pm$? To properly address this question, one should realize that the $\mathcal{R}_p$ interactions in Eq. (2) violate lepton number. Therefore, $H^\pm$ and $\tilde{\tau}^\pm$ cannot be distinguishable in terms of the SM quantum numbers. One would need to specify the complete theory and work out the allowed state mixing [19]. In this sense, the Higgs fields lose the identity as that in R -conserving SUSY theories.

In conclusion, we have studied the top-quark decays $t \rightarrow \tilde{\tau} b$ and $t \rightarrow \tau b \tilde{\chi}_1^0$ via R -parity violating interactions in SUSY theories. We found that the existing Tevatron data can put a 95% confidence level upper bound on the coupling to be 0.94 (0.63) for a long-lived (short-lived) $\tilde{\chi}_1^0$ with $m_{\tilde{\tau}} = 70$ GeV. At Tevatron Run II with an integrated luminosity of 2 (10) fb^{-1} , one can obtain a 2σ constraint as 0.38 (0.24) for a long-lived $\tilde{\chi}_1^0$ and 0.29 (0.19) for a short-lived $\tilde{\chi}_1^0$, beyond the current indirect limit. These limits from the direct search are largely independent of other SUSY parameters.

Note added: After the submission of this paper, we received an article [20] in which hadronic observables of Z decays were studied and new constraint on $|\lambda'_{333}|$ was obtained.

ACKNOWLEDGMENTS

We thank J. Feng, S. Mrenna, W. Porod and J.-M. Yang for discussions on the related topics. This work was supported in part by a DOE grant No. DE-FG02-95ER40896, in

part by the Wisconsin Alumni Research Foundation and in part by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP).

REFERENCES

- [1] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 357 (1997).
- [2] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. Lett. **82**, 4975 (1999).
- [3] J. Sender, Phys. Rev. **D54**, 3271 (1996); M. Hosch, R. J. Oakes, K. Whisnant, J. M. Yang, B.-L. Young and X. Zhang, Phys. Rev. **D58**, 034002 (1998).
- [4] C. Aulak and R. Mohapatra, Phys. Lett. **B119**, 136 (1983); F. Zwirner, *ibid.* **B132**, 103 (1983); L. J. Hall and M. Suzuki, Nucl. Phys. **B231**, 419 (1984); I. H. Lee, *ibid.* **B246**, 120 (1984); J. Ellis *et al.*, Phys. Lett. **B150**, 142 (1985); G. G. Ross and J. W. F. Valle, *ibid.* **B151**, 375 (1985); S. Dawson, Nucl. Phys. **B261**, 297 (1985); R. Barbieri and A. Masiero, *ibid.* **B267**, 679 (1986).
- [5] V. Barger, G. Giudice and T. Han, Phys. Rev. **D40**, 2987, (1989); H. Dreiner, hep-ph/9707435, published in “Perspectives on Supersymmetry”, edited by G.L. Kane, World Scientific.
- [6] H. Dreiner and R. Phillips, Nucl. Phys. **B367**, 591 (1991).
- [7] For a recent review on R -parity violation at the Tevatron, see B. Allanach *et al.*, hep-ph/9906224.
- [8] A. Bartl, W. Porod, M. A. García-Jareño, M. B. Magro, J. W. F. Valle, and W. Majerotto, Phys. Lett. **B384**, 151 (1996); L. Navarro, W. Porod and J. W. F. Valle, Phys. Lett. **B459**, 615 (1999); F. de Campos, M. A. Diaz, O. J. P. Éboli, M. B. Magro, L. Navarro, W. Porod, D. A. Restrepo and J. W. F. Valle, hep-ph/9903245.
- [9] J.-M. Yang, hep-ph/9905486; O. Lebedev, W. Loinaz and T. Takeuchi, hep-ph/9910435.
- [10] G. Bhattacharyya, D. Choudhury and K. Sridhar, Phys. Lett. **B355**, 193 (1995); G. Bhattacharyya, J. Ellis and K. Sridhar, Mod. Phys. **A10**, 1583 (1995).

- [11] J. Erler, J.L. Feng and N. Polonsky, Phys. Rev. Lett. **78**, 3063 (1997).
- [12] H. Haber and G. Kane, Phys. Rep. **117**, 75 (1985).
- [13] S. Abachi *et al.* (D0 collaboration), Phys. Rev. Lett. **79**, 1203 (1997); F. Abe *et al.* (CDF collaboration), Phys. Rev. Lett. **80**, 2779 (1998).
- [14] K. Agashe and M. Graesser, Phys. Rev. **D54**, 4445 (1996).
- [15] M. Drees, Phys. Rev. **D33**, 1468 (1986); S. Dimopoulos and G. Giudice, Phys. Lett. **B357**, 573 (1995); A. Pomarol, D. Tommasini, Nucl. Phys. **B466**, 3 (1996); J. Bagger, J. Feng, N. Polonsky, R.-J. Zhang, hep-ph/9911255.
- [16] DELPHI Collaboration, P. Abreu *et al.*, Eur. Phys. J. **C7** 595 (1999).
- [17] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **79**, 3585 (1997).
- [18] F. Borzumati, J.-L. Kneur and Nir Polonsky, Phys. Rev. **D60**, 115011 (1999).
- [19] C.-H. Chang and T.-F. Feng, hep-ph/9908295.
- [20] O. Lebedev, W. Loinaz and T. Takeuchi, hep-ph/9911479.